

Name: _____ Answer Key _____

Disease 101 Workshop Quiz

1. What is the recommended treatment for a person diagnosed with gonorrhea?
 - a. Cefixime 400 mg orally plus Azithromycin 1g Orally
 - b. Ceftriaxone 125 mg IM in a single dose plus Azithromycin 1g Orally
 - c. Ceftriaxone 250 mg IM in a single dose
 - d. Ceftriaxone 250 mg IM in a single dose plus Azithromycin 1g Orally
 2. What is the window period for HIV & Hepatitis C?
 - a. 1 month
 - b. 6 weeks
 - c. 6 months
 - d. 3 months
 3. HCV is primarily transmitted by:
 - a. Sexual activity
 - b. Tattoos
 - c. Blood
 4. An individual diagnosed with latent TB is infectious and can transmit TB to others.
 - a. True
 - b. False
 5. Which group of persons diagnosed with chlamydia are recommended to have a test-of-cure?
 - a. Pregnant women
 - b. People with multiple sex partners
 - c. Men who have sex with men
 6. Chlamydia or gonorrhea infected men and women should be retested how soon after treatment?
 - a. 2 weeks
 - b. 1 month
 - c. 3 months
 - d. 6 months
 7. Most HIV cases in North Dakota are men who have sex with men.
 - a. True;
 - b. False.
 8. HCV is the leading cause for liver transplants in the U.S.
 - a. True;
 - b. False
 9. Which statement is true?
 - a. The minority of people become chronically infected with HCV
 - b. The majority of people will clear HCV and not have ongoing disease
 - c. The majority of people who contract HCV will become chronically infected
-

10. Common symptoms of TB include:
- a. Cough
 - b. Fever
 - c. Unexplained weight loss
 - d. All of the above
11. An individual on HIV treatment is no longer infectious towards others.
- a. True
 - b. False
12. What are the correct stages of syphilis infection?
- a. Primary, Secondary, Tertiary
 - b. Primary, Secondary, Neurosyphilis
 - c. Primary, Secondary, Congenital
13. For a pregnant woman, an alternate for penicillin can be used to treat a syphilis infection.
- a. True;
 - b. False.
14. What is the most common symptom of primary syphilis?
- a. Rash
 - b. Alopecia
 - c. Chancre
 - d. Swollen lymph nodes
15. What measure is effective for decreasing the risk of reinfecting an index patient?
- a. Timely treatment of all sex partners
 - b. Treatment of current sex partner only
 - c. Referring partners in for treatment
16. The infectious period for someone diagnosed with active TB is how many months prior to symptom onset:
- a. 1 month
 - b. 3 months
 - c. 6 months
 - d. 12 months
17. Expedited partner therapy can be utilized to treat cases and contacts in North Dakota.
- a. True;
 - b. False.
18. What is the type of questions that should be asked in a counseling session?
- a. Open-ended
 - b. Closed-ended
-

19. When eliciting partner information, the following information should be collected:
- a. Name
 - b. First, Last & Frequency of Exposure
 - c. Presence of Symptoms
 - d. Locating Information – Address, Phone
 - e. All of Above
20. Which test is used to confirm active HCV infection?
- a. HCV antibody test;
 - b. HCV viral load test;
 - c. ALT test;
 - d. None of the above
-